## (b) Claims

Kindly amend claims 9 and 12 as follows.

The following is a complete listing of the claims, and replaces all earlier versions and listings.

 (Previously Presented) A magnetic toner comprising magnetic toner particles each comprising at least a binder resin and a magnetic iron oxide, wherein:

the magnetic toner has a saturation magnetization  $\sigma s$  being in the range of 5 to 60 Am<sup>2</sup>/kg and a remanent magnetization  $\sigma r$  being in the range of 0.1 to 10.0 Am<sup>2</sup>/kg in a measured magnetic field of 795.8 kA/m;

the binder resin contains a polyester component polymerized by using a Ti chelate compound having a ligand selected from the group consisting of a diol, a dicarboxylic acid, and an oxycarboxylic acid as a catalyst; and the Ti chelate compound is represented by any one of the following formulae (I) to (IV) and hydrates thereof:

Formula (I)

in the formula (I), R<sub>1</sub> denotes one of an alkylene group or an alkenylene group each having 2 to 10 carbon atoms and may have a substituent, M denotes a countercation, m denotes a cation number, n denotes a cation valence, n=2 when m=1, n=1 when m=2, and M denotes one of a hydrogen ion, an alkali metal ion, an ammonium ion, or an organic ammonium ion when n=1, or denotes an alkali earth metal ion when n=2;

Formula (II)

in the formula (II),  $R_2$  denotes one of an alkylene group or an alkenylene group each having 1 to 10 carbon atoms and may have a substituent, M denotes a countercation, m denotes a cation number, n denotes a cation valence, n=2 when m=1, n=1 when m=2, and M denotes one of a hydrogen ion, an alkali metal ion, an ammonium ion, or an organic ammonium ion when n=1, or denotes an alkali earth metal ion when n=2;

Formula (III)

in the formula (III), M denotes a countercation, m denotes a cation number, n denotes a cation valence, n=2 when m=1, n=1 when m=2, and M denotes one of a hydrogen ion, an alkali metal ion, an ammonium ion, or an organic ammonium ion when n=1, or denotes an alkali earth metal ion when n=2;

Formula (IV)

in the formula (IV),  $R_3$  denotes one of an alkylene group or an alkenylene group each having 1 to 10 carbon atoms and may have a substituent, M denotes a countercation, m denotes a cation number, n denotes a cation valence, n=2 when m=1, n=1 when m=2, and M denotes one of a hydrogen ion, an alkali metal ion, an ammonium ion, or an organic ammonium ion when n=1, or denotes an alkali earth metal ion when n=2.

- 2. (Cancelled)
- (Cancelled)
- (Original) A magnetic toner according to claim 1, wherein the magnetic iron oxide comprises 0.1 to 2.0% by mass of an Si element.
- (Original) A magnetic toner according to claim 1, further comprising hydrophobic silica treated with hexamethyldisilazane and with silicone oil.
- (Previously Presented) A magnetic toner according to claim 1,
   wherein an average circularity of the magnetic toner particles of the magnetic toner which

have equivalent circle diameters of 3  $\mu$ m or more and 400  $\mu$ m or less measured with a flow particle image analyzer, is 0.930 or more and less than 0.970.

## 7. (Cancelled)

- 8. (Previously Presented) A magnetic toner according to claim 1, wherein the polyester component comprises a compound having a structure containing oxyalkylene ether of a novolak phenolic resin as an alcohol component.
- (Currently Amended) A magnetic toner according to claim 1, further comprising a metal compound of aromatic hydroxyl carboxylic acid represented by the following formula (13)[[.]];
   Formula (13)

$$\left[ \begin{array}{c} H_2O & O \\ (B)-Z & O-C \\ C-O & Z-(B) \\ O & H_2O \end{array} \right]^{-} A^{\cdot *}$$

wherein M represents a coordinating central metal; (B) represents (i) a group of the following structure:

which may contain a substituent, wherein X represents a hydrogen atom, a halogen atom, or a nitro group; or (ii)

wherein, R represents a hydrogen atom, an alkyl group having 1 to 18 carbon atoms, or an alkenyl group having 2 to 18 carbon atoms,

A'\* represents hydrogen, a sodium ion, a potassium ion, an ammonium ion, or an aliphatic ammonium ion and Z represents -O- or -C(=O)-O-.

- (Previously Presented) A magnetic toner according to claim 1,
   wherein the Ti chelate compound is represented by the formula (I).
- 11. (Previously Presented) A magnetic toner according to claim 1, wherein the binder resin contains a polyester component polymerized by using Ti chelate compounds (1) and (2) together thereof:

Ti chelate compound (1)

Ti chelate compound (2)

$$\begin{pmatrix} H_{2}C & O & CH_{2} \\ H_{2}C & CH_{2} & CH_{2} \end{pmatrix}^{2} \cdot 2K$$

12. (Currently Amended) A magnetic toner according to claim [[1]] 9. further comprising Al hydroxycarboxylic compound and a monoazo iron compound, wherein the compound according to the formula (13) is an Al hydroxycarboxylic compound.

7